

# Uterine Leiomyoma in Kinshasa, the Capital of the Democratic Republic of Congo

Ingala P<sup>a\*</sup>, Lepira F<sup>b</sup>, Muhindo S<sup>c</sup>, Mputu L<sup>d</sup>

*<sup>a,c,d</sup>Department of Obstetrics and Gynecology, University of Kinshasa Hospital, Kinshasa, 00243, Democratic Republic of Congo*

*<sup>b</sup>Department of Internal Medicine, University of Kinshasa, Kinshasa, 00243, Democratic Republic of Congo*

## Abstract

The aim of the present study was to determine the particularities of Uterine Leiomyomas among Congolese in Kinshasa the capital of the Democratic Republic of Congo (DRC) in the present conditions of medical practices. A sample of 644 patients with uterine leiomyoma were selected from 6440 cases of uterine leiomyoma among 30395 patients treated in gynecology units of three medical institutions of Kinshasa (University hospital of Kinshasa, Saint Joseph hospital and Edith medical center) from January 1<sup>st</sup>, 2003 to December 31, 2012. The study is a descriptive one. The following variables were taken account: medical history [age, age at menarche, parity, education, civil state, history of UL, symptoms and body mass index (BMI)]; lifestyle (smoking, alcohol intake); ultrasounds characteristics; hysterosalpingographies characteristics, treatment, and direct cost of treatment. Statistical analysis were performed using Excel 12.0 software. Demographic, clinical, ultrasound, hysterosalpingography and treatment data were evaluated using descriptive statistics: mean, standard deviation (SD), and percentage (%) as appropriate. The frequency of uterine leiomyoma was 21, 18%. That one concern mainly patients at 35 years old or more [49, 6% (35-44years), ≥45years (20, 6%)], singles (70, 4%), null parous (59, 4%), having a high level of study (university: 54, 6%), history of UL (56, 7%), and alcohol intake (75, 5%). Hemorrhage (33, 2%) and pelvic pain (31, 6%) are the most frequent expression of those tumors. The most of those patients have excess weight (43, 1%) or obesity (46, 5%). The majority of uterine leiomyoma was corporeal (82, 9%) intramuscular (42, 4%) and their number didn't overtake five by patient (70, 8%) in majority of cases. Majoration of the uterine cavity (46, 5%) and Fallopian tubes obstructions (30, 6%) are the most frequent abnormalities in hysterosalpingography. Myomectomy is the main treatment (65, 2%). The mean of direct cost were 803\$ USA and 884\$ USA for myomectomy and hysterectomy respectively.

**Keys Words:** Uterine Leiomyoma -Particularities- Democratic Republic of Congo.

---

\* Corresponding author.

## 1. Introduction

The uterine leiomyoma (UL) is the most frequent genital tumor to women in reproductive age [1, 2, 3]. He is more frequent in black than white [4, 5, 6] and represents an important public health problem according to his incidence, prevalence, morbidity and cost of his management [7, 8].

In the Democratic Republic of Congo (DRC) the characteristics of UL have been explored many years ago when ultrasound was not used commonly as one of diagnosis methods [9, 10].

The objective of this paper is to evaluate the particularities of UL in the DRC regarding the present conditions of medical practices.

## 2. Materials and Methods

The study protocol was approved by the Ethical Committee of Kinshasa School of Public Health. University of Kinshasa (ESP/CE/028/2013).

6440 patients with UL have been collected in three medical institutions (University of Kinshasa hospital, Saint Joseph hospital and Edith medical center) of Kinshasa the capital of the DRC among 30395 patients who have been treated in the gynecology units from January 01, 2003 to December 31, 2012. A sample of 644 patients as been selected by random from those patients for the study.

The study is a descriptive one. Her objective was to determine the actuals particularities of UL among Congolese women in Kinshasa.

The following variables were taken account: medical history [age, age at menarche, parity, education, civil state, history of UL, symptoms and body mass index (BMI)]; lifestyle (smoking, alcohol intake); ultrasounds characteristics; hysterosalpingographies characteristics, treatment, and direct cost of treatment.

Statistical analyses were performed using Excel 12.0 software. Demographic, clinical, ultrasound, hysterosalpingography and treatment data were evaluated using descriptive statistics: mean, standard deviation (SD), and percentage (%) as appropriate.

## 3. Results

**Table I:** Frequency of uterine leiomyoma

$$F = 6440 / 30394 \cdot 100 = 21,18 \%$$

The table below show that the majority of patients have 35 years old or more [49, 6% (35-44years),  $\geq 45$ years (20, 6%)], have had their menarche at 12 year or late (71, 8%), have none parity (null parous) (59,4%) or more (multiparous) (40,6%), have reached university (54, 6%) and are single (70, 4%). History of UL (56, 7%), and alcohol intake are the main characteristics of their medical history.

Hemorrhage (33, 2%) and pelvic pain (31, 6%) are the most frequent expression their tumor. The most of those patients have excess weight (43, 1%) or obesity (46, 5%).

**Table II:** Demographic, and clinical characteristics of patients

VARIABLES	n	%
<b>Age (years)</b>		
≤ 24	10	1,5
25-34	182	28,2
35-44	320	49,6
≥45	132	20,4
<b>Age at menarche (years)</b>		
<12	182	28,2
≥12	462	71,8
<b>Parity</b>		
0	262	59,4
≥1	382	40,6
<b>Education</b>		
Primary or less	146	22,6
Secondary	146	22,6
University	352	54,6
<b>Civil State</b>		
Married	191	29,6
Single	453	70,4
<b>UL History</b>		
Not	278	43,1
Yes	366	56,9
<b>Alcohol Intake</b>		
Not	158	24,5
Yes	486	75,5
<b>Smoking</b>		
Not	616	95,6
Yes	28	4,4
<b>Symptoms</b>		
Hemorrhage	214	33,2
Pelvic Pain	110	17,0
Infertility	204	31,6
Others	96	14,9
<b>BMI</b>		
Normal	66	10,2
Excess Weight	278	43,1
Obesity	300	46,5

From the table III the majority of uterine leiomyoma was corporeal (82, 9%), intramuscular (42, 4%) and their number didn't overtake five by patient in the most of cases (70, 8%).

**Table III:** Ultrasounds characteristics of uterine leiomyomas

Ultrasounds characteristics	N	%
<b>Localisation/ uterine subdivisions</b>		
Corporeal		
Isthme	534	82,9
Cervix	84	13,0
	26	4,0
<b>Localisation/thickness</b>		
Sub sessorial	100	15,5
Intra mural	274	42,4
Sub mucosal	192	29,8
Intra cavity	78	12,1
<b>Number</b>		
≤5	456	70,8
≥6-≤10	102	15,8
≥11	66	10,2

**Table IV:** Hysterosalpingographies characteristics of uterine leiomyomas

Hystersalpingographies characteristics	N	%
<b>uterines images</b>		
normal	332	51,5
majoration of cavity	300	46,5
two cavities	4	0,6
synechies	8	1,2
<b>tubes abnormalities</b>		
normal	408	63,3
obstruction	196	30,4
hydrosalpinx	26	4,0
phimosi	14	2,1

The table IV show that majoration of the uterine cavity (46, 5%) and Fallopian tubes obstructions (30, 6%) are the more frequent hysterosalpingographies abnormalities of uterine leiomyoma observed among patients in that study.

**Table V:** Treatment

Treatment	N	%
medical	52	8,0
Myomectomy	420	65,2
Hysterectomy	172	26,7

The table V shows that myomectomy is the main treatment of UL (65, 2%).

**Table VI:** Direct cost of treatment

Medical institution	Direct cost of myomectomy	Direct cost of hysterectomy
university hospital	750\$ usa	850\$ usa
Saint joseph hospital	360\$ usa	452\$ usa
Edith medical center	1300\$ usa	1350\$ usa
<b>Mean</b>	<b>803\$usa</b>	<b>884\$usa</b>

The table VI show that the mean of direct cost of uterine leiomyoma are 803\$ USA for myomectomy and 884\$ USA for hysterectomy.

#### 4. Discussion

The present study shows that the frequency of UL is 21, 18%. Although that high level that result is comparable to those others much Africans countries [11, 12]. On contrary of Africans countries the frequencies of UL in Europeans countries are less [13, 14]. Those observations support the opinion which consider that UL is the tumor of black women. The most of patients with UL have 35 years old or more. That result join the fact of advanced age and increasing age are two parameters which are always correlated with the rising of UL frequency. For explaining that phenomenon Rongieres & al [15], and Adama & al [16] think that by growing up, UL become symptomatic toward 40 years old and their diagnosis easiest. Majority of patients are nulliparous (59,4%). That result is an argument which support the fact of a reverse relation exists between increasing of parity and occurring of UL. For many authors that relation can be understood by hormonal and no hormonal change of deliveries. Among others the reduction of number of menses cycles, the regression of the level of estrogen and progesterone, and the increasing of sex binding globulin (SBG) [17, 18]. The presence of history of UL to much patients suggest the rule of heredity in the pathogenesis of UL. Alcohol intake, excess weight and obesity are others factors which are more found to patients in that study. Those results join many others results [19, 20] and can be explained by tumor genesis action of acetaldehyde providing from the metabolism of ethanol [21] and the contribution of lipid metabolism in the production of estrogen [22]. Hemorrhage and pelvic pain as the main expression of UL is not an exception for the study [23, 24]. The most frequent localization of Uterine leiomyomas was the corporeal one (82, 9%) as for Algeria women [25]. Fallopian tubes obstructions appear as the main abnormality of UL in hysterosalpingography. That result join others studies which demonstrated that fallopian obstructions is the commonest pathology found on HSG in women presenting infertility in Kinshasa (DRC) and Kampala (UGANDA) [26, 27].

Even if hysterectomy is the final treatment of UL [28, 29], the main surgical treatment is myomectomy in the present study. We think that that result can be understood according to the high need of maternity which characterize African area.

The mean of direct cost of UL are 803\$ USA and 884\$ USA for myomectomy and hysterectomy respectively. Cravello & al [30], in France estimated direct cost to 4600FF for surgical hysteroscopy and to 7900 FF for vaginal hysterectomy. Soliman & al. [31] estimated between 11.717 and 25.023 \$ USA all expenses of UL management by patient in USA. Cardoso & al. [32] evaluated expenses of UL between 5, 9-34, 4 billion \$ USA by year in USA. The comparison of those cost evaluations suggest that the treatment of UL is less expensive in our area. But by taking account of level of life of the population, UL stay also an important problem of public health in DRC according to the cost of treatment.

## **5. Conclusion**

The present study show that the frequency of UL is 21, 18%. The patients concerned are essentially those one of 35 years old and more, null parous, singles having a high level of education, a history of UL, alcohol intake and body mass index over than the normal ranch. The majority of uterine leiomyoma are corporeal, intramuscular and their number didn't overtake five by patient in majority of cases. Majoration of the uterine cavity and Fallopian tubes obstructions are the most frequent abnormalities in hysterosalpingography. Hemorrhage and pelvic pain are the most frequent expression of those tumors. Myomectomy is the main surgical treatment. The direct cost is important for a third world country as DRC: 803\$ USA and 884\$USA for hysterectomy respectively.

## **Acknowledgement**

The authors gratefully thank, medical team of University clinics of Kinshasa, Saint Joseph Hospital and Edith medical Center for their invaluable help for the collection data of the study.

## **References**

- [1]. Sparic R., Mirkovic L., Malvasi A., Tinelli A. Epidemiology of uterine myomas: A Review Int J Fertil Steril. 2016; 9:424-35.
- [2]. Flake GP., Andersen J., Dixo D. Etiology and Pathogenesis of uterine leiomyoma: a review. Environ Health Perfect. 2003; 11:1037-1054.
- [3]. Baird DD., Dunson DB., Hill RC. Cousins D., Schectman JM. High cumulative incidence of Uterine Leiomyoma in black and white women: ultrasound evidence. Am J Obstet Gynecol. 2003 Jan; 188(1):100-7.
- [4]. Catherino WH., Eltoukhi HM., Al-Hendy. Racial and ethnic differences in the pathogenesis and clinical manifestation of uterine leiomyoma. Semin Reprod Med. 2013; 31151:370-379

- [5]. Marshall LM., Spiegelman D., Barbieri RL. Goldman ME., Manson JE et al. Variation in the incidence of uterine leiomyoma among premenopausal women by age and race. *Obstet Gynecol.* 1997; 90: 967-973.
- [6]. Parker W.H. "Etiology, symptomatology, and diagnosis of uterine myomas". *Fertility and Sterility.* 2007; .87:725-736.
- [7]. Heba M. Eltoukhi., Monica N. Modi., Meredith Weston., Alicia Y.Amonstrong., Elizabeth A.Stewart. The Health Disparities of Uterine Fibroids for African Women: A Public Health Issue. *Am J Obstet Gynecol.*2014 Mar; 210(3):194-199.
- [8]. Lauren A Wise., Julie Palmer., Elizabeth A Stewart., Lynn Rosenberg. Age-Specific Incidence Rates for Self-Reported Uterine Leiomyomata in the Black Women's Health Study. *Obstet Gynecol.* 2005 Mar; 105 (3): 563-568.
- [9]. Muberuka. Devenir des femmes zaïroises après myomectomie. Mémoire de fin de spécialisation en Médecine. Faculté de Médecine Université de Kinshasa 1979.
- [10]. Mundy M. Devenir gynécologique et obstétrical après myomectomie aux cliniques universitaires de Kinshasa. Mémoire de fin de spécialisation. Faculté de médecine.Université de Kinshasa, 1984.
- [11]. Dia A., Beye SB., Dangou JM., Dieng M., Woto gaye., Toure CT. Uterine Leiomyoma at the surgical department of the Teaching Hospital of Dakar. Report of 140 cases operated in two years. *Dakar Med.* 2003; 48(2):72-6.
- [12].Obed JY., Bakob B., Usman SD., Moruppa JY., Kadas S. Uterine Fibroids: risk of occurrence after myomectomy in Nigeria population. *Arch Gynecol Obstet.* 2011 Feb; 283(2):311-5.
- [13]. Fernandez H., Buffet CN., Allouche S. Prevalence of Uterine Fibroid in France and impact in quality of life : results of survey among 2500women between 30-55years. *J Gyneco Obst Biol Reprod,* 2014 Nov; 43(9):721-7.
- [14]. Downes E., Sikri A., Gilabert-Estelles J., Rolge SC., Dodd SL., Maroulis C., Subramanian D. *Eur J Obstet Gynecol Reprod Biol.* 2010 Sep;152(1):96-102.
- [15]. Rongières C. Recommandations pour la pratique Clinique: Epidémiologie du fibrome, facteurs de risque et fréquence. Impact en santé publique. *Gynécol Obstet Biol Reprod* 1999; 28:701-706.
- [16]. Adama Hongebila A, Modji S, Aboubakari A, Wonegouk, Akpadza K. Aspects epidemiologiques et chirurgicaux de fibromes uterins opérés à l'hôpital du district de Lomé. *J rech Sci Univer Lomé* 2012 ; serie D ; 14(1) :131-137.
- [17]. Windham GC, Elkin E, Fenster L, et al. Ovarian hormones in premenopausal women: variation by

- demographic, reproductive and menstrual cycle characteristics. *Epidemiology* 2002; 13:675-84.
- [18]. Bernstein L, Pike MC, Ross RK, et al. Estrogen and sex hormone-binding levels in nulliparous and parous women. *J Natl Cancer Inst* 1985; 74: 741-5.
- [19]. Nagata C., Nakamura K., Oba S., Hayashi M., Takeda N., Yasuda K. Association of intakes of fat, dietary fibre, soya isoflavones and alcohol with uterine fibroids in Japanese women. *Br J Nutr.* 2009 May; 101(10): 1427-31.
- [20]. Yanq Y., He Y., Zenq Q., Li S. Association of body size and body fat distribution with uterine fibroids among Chinese women. *J Womens Health (Larchmt).* 2014 Jul; 23(7): 619-26.
- [21]. Hye-Kyung Na., Ja Young Lei. Molecular Basis of Alcohol-Related Gastric and Colon Cancer. *Int J Mol Sci.* 2017 Jun; 18(6):1116. Publish online 2017 May 24. doi:10.3390/ijms18061116.
- [22]. Shikora SA., Niloff JM., Bistrian BR., Forse RA., Blackburn GL. Relationship between obesity and uterine leiomyomata. *Nutrition.* 1991 Jul-Aug; 7(4): 251-5.
- [23]. Zimmermann A., Bernuit D., Gerlinger C., Schaefer M., Geppert K. Prevalence, symptoms and management of uterine fibroids: an international internet-based survey of 21,746 women. *BMC Women Health.* 2012; 12:6-16.
- [24]. Okogbo FO, Ezechi OC, Loto OM, Ezeobi PM. Uterine leiomyomata in south western Nigeria: a clinical study of presentations and management outcome. *African Health Sciences* 2011; 2: 271-278.
- [25]. Nourelhouda Chalal., Abbassia Demmouche. Profil épidémiologique des fibromes utérins dans la région de Sidi Bel Abbès, Algeria. *Pan Afr Med J.* 2013; 15:7.
- [26]. Muhindo S. Profile of patients with uterine myomas in a population of infertile women in Kinshasa. *Memory of end of specialization.* 2013; Faculty of Medicine University of Kinshasa.
- [27]. Kiqui-Malwadde E., Byanyima RK. Structural findings at hysterosalpingigraphy in patients with infertility at two private clinics in Kampala, Uganda. *Afr Health Sci.* 2004 Dec; 4(3): 178-81.
- [28]. Nzau NE., Mboloko E., Tandu-Umba NFB., Lokengo LD. Hystérectomie aux cliniques universitaires de Kinshasa: de 2002-2010. *Médecine de l'Afrique noire.* 2012; 59:221-230.
- [29]. Gante I., Medeiros- Borges C., Aquas F. Hysterectomy in Portugal (2000-2014): what has changed? *Eur J Obstet Gynecol Reprod Biol.* 2017; 208:97-102.
- [30]. Cravello L., Isnardi M., Violin G., Duthilleul A., Sambuc R et al. Evaluation du coût direct hospitalier et extra hospitalier de l'hystéroscopie opératoire et de l'hystérectomie vaginale. *Journal de gynécologie obstétrique et biologie de la reproduction.* 1999; 28:335



- [31]. Soliman AM., Yang H., Du EX., Kellar SS., Winkel C. The direct and indirect cost of uterine fibroid tumors; a systematic review of the literature between 2000 and 2013. *Am J Obstet Gynecol.* 2015; 213:141-60.
- [32]. Cardozo ER M.D., Clark AD., MD., Nicole James H.Segars. The estimated annual cost of uterine leiomyomata in the United States. *Am J Obstet.*2012; 206:211.e1-211.e9. .

## **6. Appendix: author contributions**

PIERRE INGALA, participate in protocol elaboration, data collection and analysis and draft the manuscript

FRANCOIS LEPIRA, participate in protocol elaboration, conception and data analysis reviewed the manuscript.

SERGE MUHINDO, participate in protocol elaboration, data collection and analysis and draft the manuscript.

ARSNE MPUTU, conceived the study, participate in data analysis and reviewed manuscript.